

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

---

## “Ir (III) ION CATALYZED KINETICS AND MECHANISTIC STUDY OXIDATION OF ALANINE BY TRICHLOROISOCYANURIC ACID”

Bandana Devi shukla<sup>1</sup>, S.K. Singh<sup>1\*</sup>, H.D. Gupta<sup>2</sup>, Arshi Gautam<sup>1</sup>

<sup>1</sup>Govt. T. R.S. (Auton.) College, Rewa (M.P.) India

<sup>2</sup>Govt. Girls College, Sidhi (M.P.) India

\*Corresponding author : sanjaysinghchemtrs@gmail.com

<b>Received:</b> 15/12/2017	<b>Revised:</b> 23/12/2017	<b>Accepted:</b> 28/12/2017
-----------------------------	----------------------------	-----------------------------

### ABSTRACT

Alanine is an  $\alpha$ -amino acid that is used in the biosynthesis of proteins. It contains an  $\alpha$ -amino group, an  $\alpha$ -carboxylic acid group, and a side chain isobutyl group, making it a non-polar aliphatic amino acid. In present paper explored kinetics and Mechanistic study of oxidation of of Alanine by Trichloroisocyanuric acid which has been investigated in water-HOAc medium. The reaction is of pseudo first-order in [TCIA], fractional order in [Alanine] and first-order in [Ir (III)] ion. The velocity of the reaction increases with increase the Dielectric constant of medium and solvent polarity. Effect of temperature has been studied and thermodynamic parameters are calculated and a suitable mechanism has been suggested.

*Key words:* Alanine; Ir(III) catalyzed; fractional order; Dielectric constant of medium; thermodynamic parameters

### INTRODUCTION

The elucidation of reaction mechanism is still one of the most fascinating problems in inorganic and organic chemistry. The above study leads to work at stoichiometry, identification of intermediates and isolation of end products as an indirect support to reaction mechanism. A

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

---

number of reports on the oxidation of organic compounds by different oxidants are available in the literature<sup>5-10</sup>. In the present work explored the kinetics and mechanistic path of oxidation of Alanine by TCIA. I have chosen TCIA as an oxidant to oxidized Alanine. TCICA act an oxidant because >N-Cl bonds are polar easily undergo heterolytic fission in polar medium<sup>11</sup>.

## MATERIALS AND METHODS

The solution of TCIA (sigma-Aldrich sample) so obtained was prepared by dissolving its weighed quantity in 100% CH<sub>3</sub>COOH (B.D.H.) and kept either amber colored flask or black paper wrapped around it to save it from the action of diffused day light which alters appreciably its concentrations.

The solution of Alanine was prepared in requisite volume of glacial acetic acid. Other reagents are grade chemicals and doubly distilled water was used throughout the experiments.

## METHOD

The known volume of oxidant, acetic acid and Ir(III) ion were taken in a conical flask while substrate and rest amount of water were in another conical flask. These two-stopper flasks were placed at experimental temperature in a thermostat of sensitivity  $\pm 0.1^{\circ}\text{C}$ . After the equilibrium of the temperature, the both solutions were mixed and aliquot was withdrawn immediately and was quenched. The amount of un-reacted TCICA was estimated iodometrically with the help of standard solution of sodium thiosulphate using starch as an indicator. The titre value at zero time was taken as “a”.

The aliquots were withdrawn at regular intervals and were estimated for un-reacted TCICA. These readings are the values of (a-x) at time “t”. The experimental data were fed into the integrated form of equation for first-order reactions. The values of pseudo first-order rate constant obtained from the rate equation –

$$k = \frac{2.303}{t} \log \frac{a}{(a-x)}$$

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

Found fairly constant within the experimental error suggested that each reaction obeys first-order kinetics. The effect of [Ir (III)] on the oxidation of Alanine was determined by adding different

## RESULT AND DISCUSSION

**Effect of oxidant:** The linear plots of  $\log(a-x)$  vs. Time, suggested that the first-order rate dependency with respect to oxidant. The value of first-order rate constant evaluated from the plot is excellently in good agreement with those calculated from first-order rate equation (Fig.1) .




Fig.1: Dependence of rate on initial concentration of oxidant

**Effect of substrate:** The reaction rate increased with increase in [Alanine]. Plot of  $k_1$  versus [Alanine] initially linear passing through origin at low concentrations but at higher concentrations of substrate it bent to x-axis tends 1 to 0 orders. This confirmed the existence of equilibrium between Alanine and oxidant (TCIA) and appeared before the slow step.(Table:1)

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

Table I


[TCIA]	$\bullet$ $2.50 \times 10^{-4}$ (mol dm <sup>-3</sup> )
[Ir (III)]	$\bullet$ $1.25 \times 10^{-4}$ (mol dm <sup>-3</sup> )
HClAc-H <sub>2</sub> O	$\bullet$ 30% (v/v)
Temperature	$\bullet$ 308 K.
[Leu] $10^2$ (mol dm <sup>-3</sup> )	$10^4 k_1$ (min <sup>-1</sup> )
1.00	5.42
1.25	6.46
2.00	8.08
2.50	8.58
4.00	9.49
5.00	9.68

**Effect of [Ir(III)]:** Reaction is fully Ir(III)ion catalyzed and velocity of the reaction increases with increase the concentration of Ir(III) ion. The plot of  $k_1$  vs. [Ir(III)] ion is obtained linear with the positive unit slope, confirming that the reaction fully catalyzed.

**Effect of dielectric constant of the medium:** first-order rate constant slightly increases with increase composition of acetic acid i.e. rate slightly accelerated with increase in dielectric constant of the medium.

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research



**Fig.2: Dependence of rate on varying in concentrations of [Ir(III)]ion**

**Stoichiometry and end-product analysis:** stoichiometry of the reaction was ascertained by equilibrating the reaction mixture containing an excess of TCIA over and Ir (III), for 24 hrs. at 308K. The un-reacted oxidant (TCIA) was determined by iodometrically. The estimated amount of un-reacted TCIA showed that three moles of alanine consumes one mole of TCICA. Acetaldehyde is the identified as the end-product of the alanine-TCIA system, which characterized by the determine melting point of 2,4-dinitrophenyl hydrazone(164°C)

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

## Result and discussion

**Absence of free radical in the system:** The presence of free radicals in the system under study was tested qualitatively by addition of 1-2 ml of acrylonitrile (monomer) in about 5-6 ml of the reaction mixture employing trapping method. The non-occurrence of turbidity and white precipitate clearly indicates the absence of free radicals in the system.


Various activation parameters namely temperature coefficient, energy of activation ( $E_a$ ), frequency factor ( $A$ ), enthalpy of activation ( $\Delta H^\ddagger$ ), free energy of activation ( $\Delta G^\ddagger$ ), and entropy of activation ( $\Delta S^\ddagger$ ) for each reaction are calculated for Alanine –TCIA system and according to the reaction mechanism, rate equation and order of reaction have been discussed. (Table:2)

**Table:2**  
**Thermodynamics parameters**

Substrate	$k_1$ s <sup>-1</sup> /mol	$A$ [mol <sup>-1</sup> s <sup>-1</sup> ]	$\Delta H^\ddagger$ kJ/mol	$\Delta G^\ddagger$ kJ/mol	$\Delta S^\ddagger$ J/mol.K
<b>Alanine</b>	14.36 ±0.32	$9.02 \times 10^4$ ±0.49	68.36 ±0.37	66.69 ±0.29	183.23 ±0.36

## MECHANISM

In view of the above experimental kinetic data, facts and finding, a suitable mechanism has been proposed for the oxidation of Alanine – TCIA system as:



# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

---

Where, **S** is standing for Alanine

$$\text{Rate} = \frac{-d[\text{TCIA}]}{dt} = kK_1K_2K_3[S][\text{Br}(\text{OH})][\text{TCIA}] \dots\dots(7)$$

Equation (7) is in good agreement with the observed experimental data and results

## CONCLUSIONS

Kinetic studies utilizing TCIA as an oxidant in series of reaction lead us to conclude that the activity of it is much limited and needs to be explored in a Broadway. It possesses vital potentiality with two-electron system and displays interesting behaviors at moderate condition of temperature. The study will act as a milestone and will pave the way for future researcher to enlighten the mechanism utilizing TCIA as an oxidant for some other organic compounds like disulphide, acetophenone and substituted acetophenones, aliphatic ketones, amines and amino acids in the similar manners and also can be catalyzed by micelles like CTAB and phosphotungstic acid etc. The contribution and information through kinetic study will enrich chemical literature to a great extent in journals. Its applied aspects may be judged in lather industries, analytical, chemical separation, and identification of organic compounds and paper and pulp industries<sup>12-16</sup>.

# Electronic Journal of Advanced Research

An International Peer review E-Journal of Advanced Research

---

## REFERENCES

1. Filler R.; *Chem Rev.* 60, 21 (1965).
2. Fieser L & Fieser M.; *Reagents for organic synthesis*, 11 (Jhon Wiley, NY) 426 (1969)
3. Ot Siji Y. Kurida T. & Imoto N.; *Bull Chem.Soc Perkin II* , 949 (1963).
4. Vasudevan K.S., Venkat Subramanian N.; *Indian J. Chem.* 24 A: 304 (1985)
5. Kamble D.L., Nandibewoor S.T., *J. Phys. Org. Chem.* 11, 171–176(1998).
6. Menakshisundaram S.P., Sathiyendrian V.; *J. Chem. Res. (S)* 8 (2000) 458–468.
7. Singh H.S., Singh R.K., Singh S.M., Sisodla A.K., *J. Phys. Chem.* 81, 1044–1059,(1997).
8. Puttaswamy, Jagadeesh R.V.; *Appl. Catal. A: Gen.* 292, 259–270(2005).
9. Morris J.C., SGlyzar J.R., Winemann M.A., *J. Am. Chem. Soc.* 70 2036–2047(1948).
10. Kikkeri N. Mohana and Paanemangalore M. Ramdas Bhandarkar: *J. Chinese Chem. Soc.*, 2007, 54, 1223-1232
11. Sanju Patel: M.Phil Dissertation(2012), central library, A.P.S. University, Rewa (M.P.)
12. Frederick George Mann and Bernard Charles Saunders: “*PRACTICAL ORGANIC CHEMISTRY*” LONGMAN GROUP LIMITED London (1960) ,pp .245.
13. Frederick George Mann and Bernard Charles Saunders: “*PRACTICAL INORGANIC CHEMISTRY*” LONGMAN GROUP LIMITED London (1980) ,pp .179.
14. Atkins P., Overton T., Rourke J., Weller M. & Armstrong F.: *Shriver and Atkins: Inorganic Chemistry*, Oxford University Press (2006).pp-436
15. Priya V., Balasubramaniyan M. and Mathiyalagan N.: *J. Chem. Pharm. Res.*, 2011, 3(1):522-528
16. Patwari S.B., Khansole S.V. and Vibhute Y.B.: *J. Iran. Chem. Soc.*, Vol. 6, No. 2, 2009, pp. 399-404.